

# Title of the paper (Ariel Narrow, 26pt, Bold, Left, Sentence case)

**First\_author\_first\_name First\_author\_family\_name<sup>1#</sup>,  
Second\_author\_first\_name Second\_author\_family\_name<sup>2</sup>,  
Third\_author\_first\_name Third\_author\_family\_name<sup>3</sup>** (Ariel Narrow, 16pt, Bold, Right)

<sup>1</sup> Affiliation of author, Name of Department, Name of University, City, COUNTRY. (Ariel Narrow, 8pt)  
<sup>2</sup> Vibration and Sound Research Group (e-VIBS), Faculty of Science and Natural Resources, Universiti Malaysia Sabah, Jalan UMS, 88400 Kota Kinabalu, Sabah, MALAYSIA.  
<sup>3</sup> Faculty of Sustainable Agriculture, Universiti Malaysia Sabah, Sandakan Campus, Mile 10, Sg. Batang, 90000, Sandakan, Sabah, MALAYSIA.  
#Corresponding author. E-Mail: corresponding author's email; Tel: +6088-320000; Fax: +6088-435324.  
(Please include email, Tel and Fax number only for the corresponding author)

**ABSTRACT** *Ganoderma boninense* is a basidiomycetes fungus that causes basal stem rot disease (BSR) in oil palm trees. In Malaysia alone, the loss caused by this disease was estimated between RM 225 Million to RM 1.5 Billion in 2011 by Malaysian Palm Oil Board. Unfortunately, many planters do not realize that their fields were infected with BSR until it is too late. Several methods have been proposed for early detection of *Ganoderma boninense* infection. In this paper, Fourier transform infrared spectroscopy (FTIR) is investigated as a tool to detect the presence of *Ganoderma boninense* in oil palm tree.....(Ariel Narrow, 11pt, Justify, Single Spacing).

**KEYWORDS:** Watercress; Gluconasturtiin; PEITC; Myrosinase activity; ... (Please provide 5 suitable keywords)

Received xxxxxx Revised xxxxxx Accepted xxxxxx In press xxxxxx Online xxxxxx

© Transactions on Science and Technology

Original Article / Review Article / Data/plot With Detail Description / New Method/Instrumentation / Negative Results With Critical Argument / Concept/Design Study / New Addition / First Records / Short Communication

(Please use Palatino Linotype, 11pt font size for the rest of the paper with single spacing)

## INTRODUCTION (headings should be written in UPPER-CASE)

Oil palm (*Elaeis guineensis* Jacq.) has been known as a truly “golden crop of Malaysia” since it generated profitable export earnings for the country and truly nature’s gifts for alleviating poverty in Malaysia (Basiron - family name, 2007 - for single author; Alexander & Chong, 2013 - for 2 authors; Yunus & Radhakhrisnan, 2004). However, the oil palm industry is being jeopardized with one major disease known as Basal Stem Rot (BSR) which is mainly caused by *Ganoderma boninense* (MPOB, 2012). This disease causes serious threat to the oil palm industry in the Southeast Asian countries, especially Malaysia and Indonesia (Joy *et al.*, 2001 - use *et al.* for more than 2 authors), as which are of the major producers and exporters of palm oil in the world. Linares (2011) on the other hand found that ... However, according to Corley and Tinker (2003) infection of this disease can cause numerous yield losses and ultimately result in the destruction of basal tissues hence death of disease palms.....(please provide single enter spacing between paragraph)

This template is the official guidelines for authors of Transactions on Science and Technology journal. These guidelines include complete descriptions of the fonts, spacing, paragraphs and other related information to prepare your manuscript for publishing in Transactions on Science and Technology journal. Please do not change or adjust the margins, line spacing and other related styles in this template.

Glucosinolates (GLS) are sulphur-containing secondary metabolites found largely in cruciferous vegetables. Certain GLS are precursors to health-promoting isothiocyanates (ITCs). Watercress (*Nasturtium officinale*) contains phenyl ethyl GLS (PEGLS) or its common name, gluconasturtiin (Williams *et al.*, 2009). The hydrolysis product of PEGLS, i.e. phenyl ethyl ITC (PEITC), is proven to restrain the growth of cancer cells (Gill *et al.*, 2007). Gupta *et al.* (2014) have published a comprehensive review on the anti-cancer effects of PEITC. However, the formation of PEITC is

easily affected by various factors such as temperature, pH and presence of additives (Eylen *et al.*, 2008). PEGLS is hydrolyzed into PEITC by the naturally-occurring enzyme myrosinase in plants. This aspect needs to be investigated further because food preparation commonly involves cutting, heating and addition of other additives which may affect the PEITC formation. Currently, there are still scarce reports on the dynamic of hydrolysis of PEGLS in watercress under various external factors. Thus, this paper described the effects of temperature and pH on myrosinase activity and PEGLS hydrolysis products in watercress.

## BACKGROUND THEORY

### The Beer-Lambert Law (First level subheading - in title-case and bold)

Protocorm proliferation and regeneration were investigated on KC medium (Knudson C, 1946) supplemented with 2% (w/v) sucrose and treated with organic additives or plant growth regulators. Four types of organic additives tested are coconut water, tomato juice (10%, 15% and 20% v/v), banana pulp (25, 75 and 125 g/L) and peptone (2 g/L). Plant growth regulators tested in this study are Naphthalene acetic acid (NAA), Zeatin and 6-Benzylaminopurine (BAP) at concentrations of 2, 4, 6  $\mu\text{M}$ , respectively. Basal medium devoid of any organic additive or plant growth regulator served as control. The medium pH was adjusted to  $5.3 \pm 0.02$  and solidified with 0.8% (w/v) agar (Sigma) prior to autoclaving for 20 min at 15 psi,  $121^\circ\text{C}$ . The cultures were maintained at  $24 \pm 2^\circ\text{C}$  under a 24 h  $\text{d}^{-1}$  photoperiod with a PPF of 20–50  $\mu\text{mol m}^{-2}\text{s}^{-1}$  provided by cool white, fluorescent tubes (Philips, Malaysia).

The following (Equation 1) is an example how equation is written given as:

$$A_{\lambda} = \log(1 / R_{\lambda}) = c \varepsilon_{\lambda} I \quad (1)$$

where  $A_{\lambda}$  is absorption,  $R_{\lambda}$  is reflection,  $c$  is the concentration of the ingredient,  $\varepsilon_{\lambda}$  is extinction coefficient of the ingredient for wavelength  $\lambda$ , and  $I$  is the pathlength of the light through the sample. **All equations must be cited and/or mentioned in the main body text.**

## METHODOLOGY

### Sample Collection (First level subheading - in title-case and bold)

Healthy and infected tissues samples were collected from oil palm plantation .....

#### *Sampling site and duration* (Maximum second level subheading written in *Italic* and in sentence-case)

Healthy and infected tissues samples were collected from oil palm plantation in Sandakan, Sabah, Malaysia. Collection of trunk tissues was carried out following the method described by (Chong, 2012).

#### *Sampling transportation and preservation*


To maintain the sample conditions, they were transported in a container with vibration isolator (Chong, 2012).

**Preparation of Buffered Water and Bligh-Dyer (First level subheading - in title-case and bold)**

300 ml of double distilled water was transferred into a separating funnel and 2.04 g potassium dihydrogen phosphate ( $\text{KH}_2\text{PO}_4$ ) was added to create a 0.05 M solution. The pH was adjusted to pH 7.2 by addition of sodium hydroxide (NaOH) pellets and the mixture was extracted with 3 x 50 ml dichloromethane (DCM). The Bligh-Dyer solvent mixture was made up using buffered water: chloroform: methanol with ratio 4:5:10, respectively.

**RESULT AND DISCUSSION**

In this result, the presence of *Ganoderma* in the infected tissue was detected with the similar peak absorbance in region..... as shown in Figure 1. Table 1 specify the wavelength of the functional group ...



**Figure 1.** FTIR spectra of *Ganoderma boninense*, healthy oil palm trunk tissues and infected oil palm trunk tissues (Alexander & Chong, 2013). (Palatino Linotype, 11pt, centred, single spacing. If the caption is more than one line, please make it justify)

**Table 1.** Functional Group of *G. boninense* mycelia and healthy oil palm trunk tissues. (Palatino Linotype, 11pt, centred, single spacing. If the caption is more than one line, please make it justify)

Wavelength (cm <sup>-1</sup> )	Functional group	
	<i>Ganoderma boninense</i>	Healthy trunk tissue
3500-3200		O-H (phenol)
3400-3200	N-H (amine)	
1650-1600	C=O (amide)	C=O (amide)
1580-1500	C=N (imine)	
1470-1450	C-H (alkane)	
1400-1390		C-O(carboxylic acid)
1250-1000	C-O-C (ether)	
1100-1000		Si-O (silicone)

**CONCLUSION**

Please prepare your paper in Microsoft Words format after carefully checked the English usage and the whole content of your paper. Also, please strictly follow the format of the journal to increase the visibility of your paper amongst researchers. Please refer to the latest published paper by the journal for further example of recent format at <http://tost.unise.org/>.

130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163

## ACKNOWLEDGEMENTS

If the research is funded, please do not forget to include the funding grant name and number/code.

## REFERENCES (Arrange the reference list in alphabetical order with numbering, in Palatino Linotype, 11pt. Provide the name of all authors and **DO NOT USE et al. IN THIS REFERENCE LIST. Please make sure all references are cited in the text.**)

- [1] Basiron, Y. 2007. Palm oil production through sustainable plantations. *European Journal of Lipid Science and Technology*, 109(3), 289-295. (Example of journal paper)
- [2] Chong, K. P. 2012. An evaluation of The Ganoderma Fungal Colonisation Using Ergosterol Analysis and Quantification. *Planter*, 88(1034), 311-319.
- [3] Corley, R. H. V. & Tinker, P. B. 2003. *The Oil Palm* (4<sup>th</sup> edition). Oxford (**city of the publisher**): Blackwell Publishing (**publisher**). (Example of book)
- [4] Joy, P. P., Thomas, J., Mathew, J. & Skaria, B. P. 2001. Medicinal Plants. In: Bose, T. K., Kabir, J., Das, P. & Joy, P. P. (Eds.). *Tropical Horticulture* (Vol. 2). Calcutta (**city of the publisher**): Naya Prokash (**publisher**). (Example of book chapter)
- [5] Linares, N. V. 2011. *Analysis of Solar Water Heating Systems in Single Family Houses-Comparison between Finnish and Spanish Situation*. MSc Thesis, Tampere University of Technology, Finland. (Example of thesis)
- [6] MPOB (Malaysian Palm Oil Board). 2012. *Summary of The Malaysian Oil Palm Industry 2012* (**title of the webpage**) (<http://bepi.mpob.gov.my/index.php/en/summary-2/824-summary-2017.html> - **copy and paste the webpage URL**). Last accessed on 2 January 2021. (Example of internet document without author)
- [7] Takahashi, D. 2016. *Southeast Asian kids are 20% more active on mobile than U.S. kids* (<https://venturebeat.com/2016/01/13/southeast-asian-kids-are-20-more-active-on-mobile-than-u-s-kids/>). Last accessed on 1 August 2021. (Example of internet document with author)
- [8] Yunus, N. M. & Radhakhrisnan, V. R. 2004. Absorption of CO<sub>2</sub> and Other Acid Gases in Aqueous Solutions of Amines - A Review. *Proceedings of the 18th Symposium of Malaysian Chemical Engineers (SOMChE)*. 29-31 August, 2013, Macau, China. pp 2-9. (Example of conference proceedings. PLEASE PROVIDE FULL NAME OF THE PROCEEDINGS TOGETHER WITH THE CONFERENCE DATE, PLACE AND PAGE NUMBER)